

AD-A126 894

GEM-DIFLUOROALLYLLITHIUM: PREPARATION BY
LITHIUM-HALOGEN EXCHANGE AND UTI. (U) MASSACHUSETTS
INST OF TECH CAMBRIDGE DEPT OF CHEMISTRY
D SEYFERTH ET AL. 23 MAR 83 TR-13


1/1

UNCLASSIFIED

F/G 7/3

NL

													END FORMED - 21 - 0101



MICROCOPY RESOLUTION TEST CHART
NATIONAL BUREAU OF STANDARDS-1963-A

ADA 126094

OFFICE OF NAVAL RESEARCH
CONTRACT NO0014-82-K-0322

Task No. NR 631-618

TECHNICAL REPORT NO. 13

GEM-DIFLUOROALLYLLITHIUM: PREPARATION BY
LITHIUM-HALOGEN EXCHANGE AND UTILIZATION
IN ORGANOSILICON AND ORGANIC SYNTHESIS

by

Dietmar Seyferth, Robert M. Simon
Dennis J. Sepelak, and Helmut A. Klein

Prepared for Publication
in the
Journal of the American Chemical Society

Massachusetts Institute of Technology
Department of Chemistry, 4-382
Cambridge, Massachusetts 02139

March 23, 1983

DTIC
ELECTE
MAR 28 1983
S D
E

Reproduction in whole or in part is permitted for any purpose
of the United States Government.

This document has been approved for public release and sale;
its distribution is unlimited.

COPY

83 03 28 041

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER 13	2. GOVT ACCESSION NO. AD A126094	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) gem-Difluoroallyllithium: Preparation by Lithium-Halogen Exchange and Utilization in Organosilicon and Organic Synthesis.		5. TYPE OF REPORT & PERIOD COVERED
		6. PERFORMING ORG. REPORT NUMBER
7. AUTHOR(s) D. Seyferth, R. M. Simon, D. J. Sepelak, and H. A. Klein		8. CONTRACT OR GRANT NUMBER(s) N00014-82-K-0322
9. PERFORMING ORGANIZATION NAME AND ADDRESS Department of Chemistry, Massachusetts Institute of Technology, Cambridge, MA 02139		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS NR 631-618
11. CONTROLLING OFFICE NAME AND ADDRESS Office of Naval Research Department of the Navy Arlington, VA 22217		12. REPORT DATE March 23, 1983
		13. NUMBER OF PAGES 33
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		15. SECURITY CLASS. (of this report) Unclassified
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) Approval for public release. Distribution unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES To be published in the Journal of the American Chemical Society		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) organolithium organofluorine organic synthesis		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) gem-Difluoroallyllithium may be generated by lithiumbromine exchange between n-butyllithium and $\text{CH}_2=\text{CHCF}_2\text{Br}$ at -95°C using an <u>in situ</u> procedure. When this $\text{Li}[\text{CF}_2\text{CHCH}_2]$ preparation is carried out in the presence of chlorosilanes, aldehydes, ketones, and esters, products of type $\text{R}_3\text{SiCF}_2\text{CH}=\text{CH}_2$, $\text{RCH}(\text{OH})\text{CF}_2\text{CH}=\text{CH}_2$, $\text{RR}'\text{C}(\text{OH})\text{CF}_2\text{CH}=\text{CH}_2$, respectively, are formed, often in good yield. The factors determining the regioselectivity in additions to $\text{C}=\text{O}$ of unsymmetrically substituted allylic lithium reagents are discussed.		

gem-Difluoroallyllithium: Preparation by Lithium-Halogen Exchange and Utilization in Organosilicon and Organic Synthesis.

Dietmar Seyferth*, Robert M. Simon, Dennis J. Sepelak, and Helmut A. Klein.

Department of Chemistry, Massachusetts Institute of Technology, Cambridge, Massachusetts 02139.

ABSTRACT


gem-Difluoroallyllithium may be generated by lithium-bromine exchange between n-butyllithium and $\text{CH}_2=\text{CHCF}_2\text{Br}$ at -95°C using an in situ procedure. When this $\text{Li}[\text{CF}_2\text{CHCH}_2]$ preparation is carried out in the presence of chlorosilanes, aldehydes, ketones and esters, products of type $\text{R}_3\text{SiCF}_2\text{CH}=\text{CH}_2$, $\text{RCH}(\text{OH})\text{CF}_2\text{CH}=\text{CH}_2$, $\text{RR}'\text{C}(\text{OH})\text{CF}_2\text{CH}=\text{CH}$, respectively, are formed, often in good yield. The factors determining the regioselectivity in additions to $\text{C}=\text{O}$ of unsymmetrically substituted allylic lithium reagents are discussed.



Accession For	
NTIS GRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A	

INTRODUCTION.

In earlier work we reported the preparation of gem-difluoroallyllithium, 1, by the transmetalation reaction (eq. 1).¹ This reagent is not stable in solution, even at -95°C,




so that conventional organolithium methodology is not applicable. However, because the Si-Cl bond of triorganosilanes reacts only slowly with organolithium reagents at low temperature, slow addition of an n-butyllithium solution to a solution containing 3,3-difluoroallyltrimethyltin as well as an excess of the triorganochlorosilane (i.e., an in situ procedure) was an alternate procedure which could be applied successfully to the synthesis of $\text{R}_3\text{SiCF}_2\text{CH}=\text{CH}_2$ compounds in good yield. This in situ procedure, however, had serious limitations. When a diorganodichlorosilane (e.g., Me_2SiCl_2) was used instead of an R_3SiCl compound, disubstitution could not be effected. The first Si-Cl bond of Me_2SiCl_2 is more reactive than the Sn-C bond of $\text{Me}_3\text{SnCH}_2\text{CH}=\text{CF}_2$, so the product which was formed when disubstitution of Me_2SiCl_2 was sought was $\text{Me}_2\text{Si}(\text{n-C}_4\text{H}_9)(\text{CF}_2\text{CH}=\text{CH}_2)$ rather than $\text{Me}_2\text{Si}(\text{CF}_2\text{CH}=\text{CH}_2)_2$. Moreover, the in situ procedure could not be applied to the synthesis of alcohols containing the CF_2CHCH_2 substituent by reaction of $\text{Li}[\text{CF}_2\text{CHCH}_2]$ with aldehydes or ketones. Under the experimental conditions, the rate of the addition of n-butyllithium to the C=O bond of the substrate was faster than its reaction with $\text{Me}_3\text{SnCH}_2\text{CH}=\text{CF}_2$ and most of the latter was recovered unchanged.

In an in situ reaction in which 3-pentanone was the carbonyl compound used, the desired product, $(C_2H_5)_2C(OH)CF_2CH=CH_2$, was obtained in only 10% yield. The tedious and cumbersome method of alternate, incremental additions could be applied to the preparation of $(C_2H_5)_2C(OH)CF_2CH=CH_2$ in 75% yield, but failed to give a good product yield when benzaldehyde was the substrate used.

The difluoroallyl group, with its two C-F bonds and its reactive C=C bond, is a potentially interesting substituent in both organic and organometallic systems. Therefore, it was of interest to improve, if possible, the procedures for the generation and utilization of gem-difluoroallyllithium.

RESULTS AND DISCUSSION.

The preparation of organolithium reagents by transmetalation, in which a suitable organolithium reagent is allowed to react with an organic derivative of another metal (usually tin, lead or mercury) (eq.2) is believed to involve nucleo-




philic displacement of R from M, a polar process.² As such, it may be expected to be relatively slow at low temperature, slower than R'Li addition to the C=O bond of aldehydes and ketones. In contrast, the lithium-halogen exchange reaction, which finds many applications in organolithium preparation (eq.3),³ according to available evidence,⁴ proceeds by an



electron transfer mechanism. As such, it should be rapid even at low temperature in ether solvents, especially in the case of polyhalomethanes.

This reasoning led us to examine $\text{CH}_2=\text{CHCF}_2\text{Br}$ as an alternate precursor of gem-difluoroallyllithium. It was hoped that the $\text{CH}_2=\text{CHCF}_2\text{Br}/\text{n-C}_4\text{H}_9\text{Li}$ reaction proceeds at a rate comparable to or even faster than that of n-butyllithium addition to the carbonyl substrate when the in situ procedure is used.

A precursor of the required halide, $\text{CH}_2=\text{CHCF}_2\text{Br}$, had been prepared by Tarrant and Lovelace by the benzoyl peroxide-induced addition of dibromodifluoromethane to ethylene in an autoclave at 80°C (eq.4).⁵ The reported runaway exotherm which resulted



in loss of most of the contents of the autoclave through the rupture disk (designed to withstand 1250 p.s.i.)⁴ was somewhat disquieting. However, these workers used an unusually large amount of benzoyl peroxide to initiate the addition⁵ and we found this reaction to be a safe and useful preparation of $\text{BrCH}_2\text{CH}_2\text{CF}_2\text{Br}$ when smaller amounts of benzoyl peroxide were used. Dehydrobromination of $\text{BrCH}_2\text{CH}_2\text{CF}_2\text{Br}$ using a saturated aqueous solution of KOH at $120\text{--}150^\circ\text{C}$ gave a ~5:2 mixture of $\text{CH}_2=\text{CHCF}_2\text{Br}$ and $\text{CF}_2=\text{CHCH}_2\text{Br}$ in yields as high as 93%. Slow distillation of this mixture through a glass helices-packed column increased the $\text{CH}_2=\text{CHCF}_2\text{Br}/\text{CF}_2=\text{CHCH}_2\text{Br}$ ratio to 20:1, and this material was used in our subsequent studies.

The lithium-halogen exchange reaction, as expected, could be applied to good advantage to the in situ generation of gem-

difluoroallyllithium. When the $\text{CH}_2=\text{CHCF}_2\text{Br}/\underline{n}\text{-C}_4\text{H}_9\text{Li}$ reaction was carried out in a 5:1:1 (by volume) mixture of THF/ Et_2O /pentane at -95°C under nitrogen in the presence of an excess of a triorganochlorosilane, good yields of the expected $\text{R}_3\text{Si}-\text{CF}_2\text{CH}=\text{CH}_2$ were obtained (Table 1). Although the difluoroallyllithium is generated by reaction at the CF_2 terminus of the difluoroallyl precursor in the $\text{CH}_2=\text{CHCF}_2\text{Br}/\underline{n}\text{-C}_4\text{H}_9\text{Li}$ reaction, vs the CH_2 terminus in the case of the $\text{Me}_3\text{SnCH}_2\text{CH}=\text{CF}_2/\underline{n}\text{-C}_4\text{H}_9\text{-Li}$ reaction, the reagent formed in these different reactions appears to be the same: $\text{R}_3\text{SiCF}_2\text{CH}=\text{CH}_2$ is formed in either case. No trace of the other isomer,⁸ $\text{R}_3\text{SiCH}_2\text{CH}=\text{CF}_2$, was observed. Noteworthy is the fact that $\text{Me}_2\text{Si}(\text{CF}_2\text{CH}=\text{CH}_2)_2$ was obtained in 73% yield when Me_2SiCl_2 was the silicon halide used. Thus, in this application, the lithium/halogen exchange synthesis of gem-difluoroallyllithium is far superior to that using the transmetalation reaction.

The in situ lithium-halogen exchange route to gem-difluoroallyllithium made possible the difluoroallylation of aldehydes and ketones as well. Dialkyl ketones (Me_2CO , Et_2CO and cyclohexanone) and an alkyl aryl ketone (acetophenone) were found to give products of type $\text{R}_2\text{C}(\text{OH})\text{CF}_2\text{CH}=\text{CH}_2$ in good to fair yield (Table 1) but no difluoroallyl product was obtained with benzophenone. Aliphatic aldehydes could be converted to alcohols of type $\text{RCH}(\text{OH})\text{CF}_2\text{CH}=\text{CH}_2$ in good yield (Table 1), but competing n-butyllithium addition became important when the $\text{C}=\text{O}$ bond was more reactive, as in the case of acrolein and benzaldehyde (Scheme 1). Thus there are some

TABLE 1. Reactions of In-Situ Generated gem-Difluoroallyl-lithium.






Reactant	Product	Yield, %
Me_3SiCl	$\text{Me}_3\text{SiCF}_2\text{CH=CH}_2$	27
Et_3SiCl	$\text{Et}_3\text{SiCF}_2\text{CH=CH}_2$	51
$n\text{-Pr}_3\text{SiCl}$	$n\text{-Pr}_3\text{SiCF}_2\text{CH=CH}_2$	50
PhMe_2SiCl	$\text{PhMe}_2\text{SiCF}_2\text{CH=CH}_2$	71
Me_2SiCl_2	$\text{Me}_2\text{Si}(\text{CF}_2\text{CH=CH}_2)_2$	74
$n\text{-C}_4\text{H}_9\text{CH=O}$	$n\text{-C}_4\text{H}_9\text{CH(OH)CF}_2\text{CH=CH}_2$	87
$(\text{CH}_3)_3\text{CH=O}$	$(\text{CH}_3)_3\text{CCH(OH)CF}_2\text{CH=CH}_2$	95
$\text{CH}_2=\text{CH-CH=O}$	$\text{CH}_2=\text{CHCH(OH)CF}_2\text{CH=CH}_2$	20
	$(\text{CH}_2=\text{CHCH(OH)C}_4\text{H}_9\text{-}n$	51)
		15
	$(\text{C}_6\text{H}_5\text{CH(OH)(CH}_2)_3\text{CH}_3$	78)
$(\text{CH}_3)_2\text{C=O}$	$(\text{CH}_3)_2\text{C(OH)CF}_2\text{CH=CH}_2$	41

TABLE 1 con't.

<u>Reactant</u>	<u>Product</u>	<u>Yield, %</u>
$(C_2H_5)_2C=O$	$(C_2H_5)_2C(OH)CF_2CH=CH_2$	70
		59
$PhC(O)CH_3$	$PhC(Me)(OH)CF_2CH=CH_2$	73
$ClCH_2CO_2Me$	$ClCH_2C(O)CF_2CH=CH_2$	95
$(CH_3)_2CHCO_2Me$	$(CH_3)_2CHC(O)CF_2CH=CH_2$	62
$(CH_3)_3CCO_2Me$	$(CH_3)_3CC(O)CF_2CH=CH_2$	49


limits on the applicability of this procedure. The C=O bond reactivity toward nucleophiles can vary widely as the substituents on the carbon atom are changed. It appears that aromatic substituents result in a C=O group reactive enough to trap most of the n-butyllithium before it can undergo the electron transfer reaction with $\text{CH}_2=\text{CH}-\text{CF}_2\text{Br}$.

gem-Difluoroallyllithium also was found to ylate esters when the in situ procedure was used (eq.5). The



yields of 1,1-difluoroallyl ketones obtained were good (Table 1).


The alkylation of esters by organolithium compounds to give ketones proceeds via initial addition of RLi to the C=O function, followed by elimination of lithium alkoxide (eq.6).



In our in situ difluoroallylation of ketones the elimination step (b) apparently does not occur at the low reaction temperature, rather during the warm-up period. This was shown in a

reaction in which it was attempted to prepare $\text{ClCH}_2\text{C}(\text{OH})(\text{CF}_2\text{CH}=\text{CH}_2)_2$ by treating methyl chloroacetate with two molar equivalents of gem-difluoroallyllithium. When this reaction mixture was treated with trimethylchlorosilane prior to warming to room temperature, the product was not the trimethylsilyl derivative of the expected carbinol, rather it was $\text{ClCH}_2\text{C}(\text{OMe})(\text{OSiMe}_3)\text{CF}_2\text{CH}=\text{CH}_2$. Thus it was the first intermediate, $\text{ClCH}_2\text{C}(\text{OMe})(\text{OLi})\text{CF}_2\text{CH}=\text{CH}_2$, which was the major species present when the trimethylchlorosilane was added.


As Table 1 shows, all products obtained in reactions of gem-difluoroallyllithium with aldehydes, ketones and esters were the ones in which the new C-C bond had been formed at the CF_2 terminus of the reagent. This observation requires some discussion. First we must consider the nature of the gem-difluoroallyllithium species. The most recent work suggests that allyllithium exists in ethereal solvents in the form of symmetrically bridged species 2,⁹ a tight ion pair




2

in which covalent contributions are important. A terminally disubstituted lithium reagent, $\text{Li}[\text{CX}_2\text{CHCH}_2]$, should have an

unsymmetrical charge distribution, 3 or 4, with the lith-



3




4

ium ion no longer bridging the terminal carbon atoms in a symmetrical fashion. Of the two extrema, 3 and 4, we expect 4 (X = F) to best describe gem-difluoroallyllithium in solution. Although the -I inductive effect of the fluorine substituents might be expected to stabilize a carbanion center, this effect will be cancelled in large part by the destabilizing repulsion between the lone pair electrons on the fluorine substituents and the electrons in the carbanion orbital.¹⁰ The work of Streitwieser and Mares¹¹ on the relative stabilizing effects of a 9-F vs a 9-CF₃ substituent in the fluorenyl anion may be noted in this connection. Also pertinent is the observation by Hine et al.¹² that one fluorine substituent on the α -carbon atom of ethyl acetate (i.e., in CFH₂CO₂Et) slightly increases the acidity of the remaining hydrogen atoms of the methyl group, but that substitution of a second fluorine atom (to give CF₂HCO₂Et) decreases the acidity of the remaining hydrogen atom by a factor of 1000 vs CH₃CO₂Et itself.

If 4 best describes gem-difluoroallyllithium, i.e., if more of the negative charge resides at the CH₂ terminus than at the CF₂ terminus,¹³ how can one reconcile this with the

observation that the products which result are those in which the new bond to the electrophile is formed at the CF₂ terminus? We suggest that the lithium counterion must be considered in any explanation which is offered. If allylic lithium reagents exist in ether solvents in the form of tight ion pairs in which there is a significant covalent bonding contribution, as indicated in 2, 3 and 4, then the lithium ion would be expected to be coordinated at the site of greatest negative charge, i.e., at the CH₂ terminus in the case of Li[CF₂CHCH₂]. This would serve to block the CH₂ terminus from attack by an electrophile relative to the "free" CF₂ terminus. An S_E2' process would result in most cases (Scheme 2). In Scheme 2 attack at a C=O group is shown, but the same considerations are applicable to attack at the Si atom of a chlorosilane.

This explanation for the observed regioselectivity of the reactions of gem-difluoroallyllithium may be applied to the reactions of other terminally substituted allylic lithium reagents. For instance, a much more complicated picture had emerged in our study of the reactions of gem-dichloroallyllithium with aldehydes and ketones.¹⁴ In the case of gem-dichloroallyllithium, we suggest that, for the same reasons, the charge density will be greater at the CH₂ terminus than at the CCl₂ end, but that this charge density difference will not be as great. It is known that chlorine atoms, while showing some effects of Cl lone pair repulsion of the negative charge at adjacent sp² carbanion centers, do so to a lesser



Scheme 2.

degree.¹⁵ As the charge density at the CH_2 terminus decreases on going from $\text{Li}[\text{CF}_2\text{CHCH}_2]$ to $\text{Li}[\text{CCl}_2\text{CHCH}_2]$, the Li^+ ion will be displaced less from the symmetrical position of unsubstituted allyllithium. Thus, the CH_2 terminus will be less blocked to attack, especially ^{to attack} by the stronger electrophiles. It is then no surprise that substrates with more reactive $\text{C}=\text{O}$ bonds such as hexafluoroacetone, 1,1,1-trifluoroacetone, trifluoromethyl phenyl ketone, benzophenone, acrolein and acetophenones with electron-withdrawing substituents on the phenyl ring react with gem-dichloroallyllithium to form the new $\text{C}-\text{C}$ bond at the CH_2 terminus. This is, after all the site of greater electron density. On the other hand, gem-dichloroallyllithium reacted with dialkyl ketones, which have a less reactive carbonyl group, to form products type $\text{R}_2\text{C}(\text{OH})-\text{CCl}_2\text{CH}=\text{CH}_2$ (after hydrolytic workup). Aliphatic aldehydes and acetophenone reacted to give mixtures of both types of products, $\text{RCH}(\text{OH})\text{CCl}_2\text{CH}=\text{CH}_2$ and $\text{RCH}(\text{OH})\text{CH}_2\text{CH}=\text{CCl}_2$, in the case of aldehydes. Unfortunately, the regioselectivity of gem-difluoroallyllithium could not be probed to the limit of the highly reactive carbonyl compounds since, as noted above, with these the addition of n-butyllithium to the $\text{C}=\text{O}$ bond was faster than its reaction with $\text{CH}_2=\text{CHCF}_2\text{Br}$. In our report on gem-dichloroallyllithium¹⁴ we attempted to rationalize the observed regioselectivity of this reagent, which obviously showed the operation of an electronic effect, in terms of the hard/soft-acid/base approach. That, however, was merely a rationalization, not an explanation. The present approach, we feel, is much

more satisfactory.


The regioselectivities of three other allylic lithium reagents which bear a chlorine substituent on one of the terminal carbon atoms also are of interest: gem-chloro-(methyl)allyllithium, $\text{Li}[\text{CH}_3\text{CClCHCH}_2]$,¹⁶ chloroallyllithium, $\text{Li}[\text{ClCHCHCH}_2]$,¹⁷ and gem-chloro(trimethylsilyl)allyllithium $\text{Li}[\text{Me}_3\text{SiCClCHCH}_2]$.¹⁸ In the first of these, the methyl group will inductively destabilize a negative charge at the carbon atom to which it is attached and the chlorine substituent will cause some destabilization as well by the lone pair effect discussed above. The observed formation of only products of type $\text{R}_2\text{C}(\text{OH})\text{C}(\text{CH}_3)(\text{Cl})\text{CH}=\text{CH}_2$ in reactions of gem-chloro(methyl)-allyllithium with cyclohexanone, methyl isopropyl ketone, acetophenone and benzophenone is easily understood on this basis. Only when steric effects become prohibitive, as in the case of methyl t-butyl ketone, was C-C bond formation at the CH_2 terminus observed.

In chloroallyllithium, destabilization by only one chlorine substituent on one of the terminal carbon atoms is operative. One might then expect to find C-C bond formation at the CH_2 terminus more favorable than in the case of $\text{Li}[\text{CCl}_2\text{CHCH}_2]$. This is what was observed.¹⁷ For instance, in the reaction of chloroallyllithium with $\text{Et}_2\text{C}=\text{O}$, the CH_2/CHCl terminus attack ratio was 3. In comparison, $\text{Li}[\text{CCl}_2\text{CHCH}_2]$ reacted with this ketone to give only $\text{Et}_2\text{C}(\text{OH})\text{CCl}_2\text{CH}=\text{CH}_2$.¹⁴ In the case of aldehydes, the formation of $\text{RCH}(\text{OH})\text{CH}_2\text{CH}=\text{CHCl}$ is highly favored, which stands in marked contrast to the regioselectivity




observed in $\text{Li}[\text{CCl}_2\text{CHCH}_2]/\text{RCHO}$ reactions.¹⁴

In the case of gem-trimethylsilyl(chloro)allyllithium the trimethylsilyl group brings two effects: the well-known stabilization of an adjacent negative charge (by $d_\pi\text{-}p_\pi$ bonding and/or by polarization effects) and a fairly substantial steric effect. Thus, while the electronic effect of the Me_3Si group would work against that of the chlorine substituent, its steric effect would hinder C-C bond formation at the substituted terminus of the allylic reagent. On reaction with 1,1,1-trifluoroacetone, only $(\text{CH}_3)(\text{CF}_3)\text{C}(\text{OH})\text{CH}_2\text{CH}=\text{C}(\text{Cl})(\text{SiMe}_3)$ was formed, but with cyclohexanone and acetophenone both possible isomeric products were obtained.

Similar, previously reported dramatic differences in regioselectivity in reactions of alkyl-substituted allylic lithium reagents vis-a-vis acetone and hexafluoroacetone¹⁹ now find a ready explanation. As noted already, an alkyl group will inductively destabilize an adjacent negative charge. That this is so in terminally alkyl-substituted allyllithium reagents has been demonstrated experimentally by ^{13}C NMR studies.²⁰ Thus, the tight ion pair found for alkylallyllithium reagents in ether solvents should be best described by 5. Barring prohibitive steric factors, formation of the new C-C bond on



reaction with aldehydes and ketones at the substituted terminus should be favored. This has been found to be generally true,²¹ and we cite only some results from our own previous work¹⁸ (eq.6-8).



As expected on the basis of the ideas developed above, all three reagents shown in eq.6-8 reacted with hexafluoroacetone to favor formation of the product alcohol with the new C-C bond at the CH₂ terminus. With n-hexylallyllithium, 68%, with Li[cyclo-C₆H₁₀CHCH₂], 100%, and with Li[EtCHC(Me)CH₂], 65%, of the product was formed at the CH₂ end of the reagent.

These considerations lead us to conclude that much remains to be done in the area of allylic lithium reagent chemistry. If the tight binding between the allyl anion and the lithium ion even in Et_2O and THF solution indeed is the crucial factor which determines the observed regioselectivities with the various electrophiles used, then the effect of strong Lewis base additives, such as $\text{Me}_2\text{NCH}_2\text{CH}_2\text{NMe}_2$, $(\text{Me}_2\text{N})_3\text{PO}$, crown ethers, cryptands, etc., which might compete in the binding of Li^+ , would be a fruitful area of study, both from the mechanistic and the synthetic point of view. Such a study might be carried out with gem-dichloroallyllithium, whose reactivity as an ambident nucleophile in the absence of additives has been well mapped.¹⁴

This research, then, has provided a new and useful route to gem-difluoroallyllithium which allows the synthesis of α , α -difluoroallyl-substituted secondary and tertiary alcohols, ketones and silicon compounds. In addition, a useful explanation of the regioselectivities observed in the reactions of all kinds of substituted allylic lithium reagents with various electrophiles has been proposed.

The introduction of fluorine in place of hydrogen in organic compounds can cause profound (and, in some cases, useful) changes in biological effects²² and the availability of the difluoroallyl anion as a reagent offers a new method of introducing fluorine into compounds of biological interest. The difluoroallylic alcohols prepared in this study may themselves be of interest as potential antimetabolites.

EXPERIMENTAL

General Comments

All reactions of gem-difluoroallyllithium were carried out under an inert atmosphere in flame-dried glassware. THF and diethyl ether were purified by distillation from sodium benzophenone ketyl under an inert atmosphere. Pentane was similarly distilled, but from LiAlH_4 . Solvents were stored under an inert atmosphere prior to use, and diethyl ether was refrigerated as well. Dimethyl ether was purified as a gas by passing it through drying columns containing Drierite, P_2O_5 , and activated alumina. It was then condensed into a flame-dried graduated cylinder at -78°C and used immediately. Chlorosilanes were purified by distillation from magnesium chips and were checked for purity by gas-liquid chromatography (GLC) before use. n-Butyllithium in hexane was purchased from the Alfa/Ventron Division of Thiokol Corporation and was used as received. All alkyllithium reagents were standardized using the method of Kofron and Baclawski.²³

Analytical GLC was performed on a Hewlett-Packard Model 5754 gas chromatograph equipped with a thermal conductivity detector. Preparative GLC was performed on either a Hewlett-Packard/F&M Model 700, Hewlett-Packard/F&M Model 720, or Gow-Mac Model 550P gas chromatograph. GLC columns were constructed from 1/4-inch o.d. copper tubing and were packed with coated diatomaceous silica supports. The silica supports were acid-washed and were treated with dimethyldichlorosilane and methanol before being coated with either General Electric Co.

SE-30 dimethylsilicone rubber gum or General Electric Co. QF-1 fluorosilicone oil.

Infrared spectra were recorded using a Perkin-Elmer Model 457A or a Perkin-Elmer Model 283B grating infrared spectrophotometer. Proton magnetic resonance spectra were recorded using a Varian Associates T-60 or a Hitachi-Perkin-Elmer R-24B spectrometer, operating at 60 MHz. High-field proton magnetic resonance spectra were recorded using a JEOL FX-90Q spectrometer, operating at 90 MHz, or a Bruker WM-250 or WM-270 spectrometer, operating at 250 MHz and 270 MHz, respectively. Chemical shifts are reported in δ units, parts per million downfield from internal tetramethylsilane.

Refractive indices were recorded using a Zeiss-Abbé type refractometer, thermostated either at 20°C or 25°C. Melting points were determined on analytically pure samples using a Büchi Capillary Melting Point Apparatus and are uncorrected.

In Situ Reaction Procedure

The standard apparatus for the reactions of gem-difluoroallyllithium generated in situ by lithium-halogen exchange consisted of a three-necked, 200 ml Morton flask equipped with an overhead mechanical stirrer, a pressure-equalizing addition funnel capped with a rubber septum, and a Claisen adapter with low-temperature thermometer and a gas-inlet adapter leading to an inert gas line (argon or prepurified nitrogen).

The apparatus was cooled and maintained at low temperatures by placing it in a wide-mouthed Dewar flask partially filled

with liquid nitrogen. By raising or lowering the flask on a laboratory jack, the reaction apparatus can be either immersed in liquid nitrogen or held just above the surface of the liquid to adjust the temperature. Low temperatures were measured using pentane thermometers (W. H. Kessler Co., Inc., +30°C to -200°C) with only the bulb immersed in the solution. Pentane total immersion thermometers read -70°C (bulb immersed) versus -78°C (total immersion) in a dry ice-acetone bath, so the temperatures reported in this thesis are probably 8-10° high for these low temperature reactions.

Preparation of $\text{BrCF}_2\text{CH}_2\text{CH}_2\text{Br}$

A high pressure stainless steel reaction vessel of 1.4 liter capacity was conditioned by performing the procedure described below using one-tenth the amount of CF_2Br_2 described. Immediately after conditioning, it was charged with 198.7 g (0.95 mol) of CF_2Br_2 and 1.13 g (4.7 mmol, 0.005 molar equivalent) of benzoyl peroxide. The bomb was assembled, mounted on a rocker, and attached to a gas line. The apparatus was charged with 120 psi ethylene (1.4 liters at 8.0 atm, compressibility factor = 0.9451 at 125 psi and 21°C,²⁴ 0.49 mol ethylene), heated to 80°C and allowed to react for 18 hours. During this time, some of the ethylene was consumed. The bomb was allowed to cool, was disassembled, and the material was emptied from the bomb and weighed (143.7 g).

This product mixture was trap-to-trap distilled (0.15 Torr, 50°C) to remove all volatiles from the brown solid residue left by the decomposition of benzoyl peroxide. The clear distillate

was again trap-to-trap distilled (atmospheric pressure, 30°C) into a flask cooled in dry ice/isopropyl alcohol slush. This was done to collect unreacted CF_2Br_2 , and the distillate weighed 74.2 g. The pot residue from this procedure was analyzed by GLC (15% SE-30 on Chromosorb W, 75°C) and showed two peaks, one for residual CF_2Br_2 and one for the expected product, $\text{BrCF}_2\text{CH}_2\text{CH}_2\text{Br}$. No higher boiling components were seen, even at elevated column temperatures. The pot residue was distilled at reduced pressure. At 160 Torr, a further 27.9 g of CF_2Br_2 distilled at 25° and was collected in a receiver cooled in a dry ice/isopropyl alcohol slush. The pressure was reduced further and the expected product, $\text{BrCF}_2\text{CH}_2\text{CH}_2\text{Br}$, n_D^{25} 1.4463 (lit.⁵ n_D^{25} 1.4450), was collected at 61-67°C, 97 Torr (literature bp = 62°C, 86 Torr).⁵ The yield was 36.7 g (154 mmol, 78%).

NMR (90 MHz, $\text{CDCl}_3/\text{CHCl}_3$): δ 2.90 [tt, $^3\text{J}(\text{H-H})$ 7.82 Hz, $^3\text{J}(\text{H-F})$ 12.5 Hz, 2H, $\text{BrCF}_2\text{-CH}_2\text{-CH}_2$], 3.47 [t, $^3\text{J}(\text{H-H})$ 7.82 Hz, 2H, $\text{CH}_2\text{-CH}_2\text{Br}$]

The yield in this reaction is based on the estimated amount of dibromodifluoromethane which reacted. This estimate was arrived at by subtracting from the total amount of CF_2Br_2 which was used, the amount of recovered CF_2Br_2 as well as the amount of CF_2Br_2 lost in handling operations, especially the venting of excess pressure from the bomb to the outdoors after the reaction is complete.

Preparation of 3,3-Difluoro-3-bromopropene

To a 500 ml round-bottomed flask containing a magnetic stirring bar was added 202.3 g (3.61 mol) of reagent grade pellets of KOH and 75 ml of distilled water. The KOH pellets dissolved partially to give a saturated aqueous solution at room temperature. To this was added 64 g (0.27 mol) of $\text{BrCF}_2\text{-CH}_2\text{CH}_2\text{Br}$, and the flask was attached to a distillation apparatus consisting of a 30 cm Vigreux column, a water-cooled distillation head, and a receiver cooled in a dry ice/isopropyl alcohol slush. An oil bath, with a magnetic stirring bar, a resistive heating coil connected to a variable transformer, and a thermometer, was placed around the reaction flask. A magnetic stirrer was placed under the oil bath to drive both the stirring bar in the bath and the one in the reaction flask.

When the temperature of the oil bath reached 120°C , a colorless liquid began to distill at temperatures ranging from $25^\circ\text{-}95^\circ\text{C}$. At the upper end of the temperature range, the distillate was cloudy and mixed with water. The temperature of the oil bath was increased slowly to 160°C to force over all of the product. After the distillation was completed, the cooled receiver flask was allowed to warm to room temperature. The dried product (Molecular Sieves, 4A) was redistilled through a 30 cm column packed with 3/16-inch glass helices. The fraction distilling from $40\text{-}45^\circ\text{C}$ was collected in a cooled (-78°C) receiver. NMR analysis of this product showed it to be a 20/1 mixture of $\text{CH}_2=\text{CH-CF}_2\text{Br}$ / $\text{CF}_2=\text{CH-CH}_2\text{Br}$. The yield was 93% (39.1 g, 0.25 mol). NMR spectral data are given in the following Figure.




Fig. 4. High-Field Proton NMR Spectrum of 3,3-Difluoro-3-bromopropene

Reactions of gem-Difluoroallyllithium with Chlorosilanes at

-95°C

A reaction with triethylchlorosilane is described in detail. Into the standard reaction apparatus containing 80 ml of THF was introduced 1.69 g (10.8 mmol) of 3,3-difluoro-3-bromopropene and 2.80 ml (16.6 mmol) of Et_3SiCl (Petrarch), followed by 20 ml of diethyl ether and 20 ml of pentane. The reaction mixture was cooled to -95°C and 7.0 ml (15.1 mmol) of a 2.2 M solution of *n*-butyllithium in hexane was added slowly, dropwise. After the addition was completed, the reaction mixture was stirred at -95°C for 90 min and then was allowed to warm slowly to room temperature. The reaction mixture was worked up in what shall henceforth be called, "the usual manner". Approximately 20 ml of distilled water was added to dissolve the lithium salts. The organic phase was drawn off using a separatory funnel, and the aqueous phase was extracted with pentane. The organic phases were combined, dried over anhydrous MgSO_4 , and concentrated using a rotary evaporator. The expected product, $\text{Et}_3\text{SiCF}_2\text{CH}=\text{CH}_2$, n^{25}_D 1.4217, was isolated and analyzed by GLC (15% SE 30 Chromosorb P, 135°C). The yield was 51%.

^1H NMR($\text{CCl}_4/\text{C}_6\text{H}_6$): δ 0.15-1.4 (m, 15H, Et_3Si) and 5.1-6.6 ppm (m, 3H, $\text{CH}=\text{CH}_2$).

IR (thin film): ν (C=C) 1645(w) cm^{-1} .

Anal. Calcd. for $\text{C}_9\text{H}_{18}\text{F}_2\text{Si}$: C, 56.21; H, 9.43.

Found: C, 56.15; H, 9.42.

The following in situ reactions were carried out in similar

manner.

- (1) gem-Difluoroallyllithium (from 19.0 mmol of $\text{CH}_2=\text{CHCF}_2\text{Br}$ and 15 mmol of n-butyllithium) reacted with 30.0 mmol of PhMe_2SiCl (Petrarch) to give $\text{PhMe}_2\text{SiCF}_2\text{CH}=\text{CH}_2$ in 71% yield. The proton NMR spectrum was in good agreement with that reported earlier.¹
- (2) gem-Difluoroallyllithium (from 10.9 mmol of $\text{CH}_2=\text{CHCF}_2\text{Br}$ and 10.8 mmol of $\text{C}_2\text{H}_5\text{Li}$ in Et_2O) reacted with 20 mmol of Me_3SiCl to give $\text{Me}_3\text{SiCF}_2\text{CH}=\text{CH}_2$ ¹ in 27% yield.
¹H NMR($\text{CCl}_4/\text{C}_6\text{H}_6$): δ 0.12 (s, 9H, Me_3Si) and 5.0-6.3 ppm (m, 3H, $\text{CH}=\text{CH}_2$).

When n-butyllithium was used to generate gem-difluoroallyllithium instead of ethyllithium the yield of $\text{Me}_3\text{SiCF}_2\text{CH}=\text{CH}_2$ was higher (89% by GLC), but the product proved to be difficult to separate by distillation from the n-butyl bromide formed in the lithium/halogen exchange reaction.

- (3) gem-Difluoroallyllithium (from 20.6 mmol of $\text{CH}_2=\text{CHCF}_2\text{Br}$ and 20.3 mmol of n-butyllithium) reacted with 10.3 mmol of Me_2SiCl_2 to give $\text{Me}_2\text{Si}(\text{CF}_2\text{CH}=\text{CH}_2)_2$, n^{25}_D 1.4016, in 74% yield. In this reaction the n-butyllithium was added in two portions: initially 10 mmol, then, after the reaction mixture had been stirred at -97°C for 25 min, another 10.3 mmol.

NMR($\text{CCl}_4/\text{C}_6\text{H}_6$): δ 0.33 (s, 6H, Me_2Si) and 5.1-6.3 ppm (m, 6H, $\text{CH}=\text{CH}_2$). IR (thin film): ν (C=C) $1640(\nu)\text{cm}^{-1}$.
 Anal. Calcd. for $\text{C}_8\text{H}_{12}\text{F}_4\text{Si}$: C, 45.27; H, 5.70. Found: C, 45.32; H, 5.63.

Reactions of gem-Difluoroallyllithium with Aldehydes and Ketones

The reaction with pentanal is described to illustrate the procedure used.

The standard reaction apparatus was charged with 80 ml of THF, 1.72 g (10.9 mmol) of $\text{CH}_2=\text{CHCF}_2\text{Br}$ and 2.35 mol (22.1 mmol) pentanal, $n\text{-C}_4\text{H}_9\text{CHO}$, 20 ml of diethyl ether and 20 ml of pentane. The solution was cooled to -95°C and 5.3 ml (12 mmol) of a 2.3M solution of n -butyllithium in hexane was added dropwise. After the addition had been completed, the reaction mixture was stirred at -95°C for 90 min and then was allowed to warm slowly to room temperature. The reaction mixture was treated with saturated aqueous NH_4Cl . The organic layer was separated and the usual work-up (as above) followed. GLC analysis of the organic liquid residue (15% SE 30 on Chromosorb P, 100°C) and collection of the product (at 135°C) showed that $n\text{-C}_4\text{H}_9\text{CH}(\text{OH})\text{CF}_2\text{CH}=\text{CH}_2$, $n^{25}\text{D}$ 1.4271, had been formed in 87% yield. NMR($\text{CCl}_4/\text{C}_6\text{H}_6$): δ 0.66-1.85, (m, 10H, $n\text{-C}_4\text{H}_9$ and OH), 3.22-3.98 (m, 1H, $\text{CH}(\text{OH})$) and 5.28-6.45 ppm (m, 3H, $\text{CH}=\text{CH}_2$). IR (film): ν (C=C) 1647(w) cm^{-1} .

Anal. Calcd. for $\text{C}_8\text{H}_{14}\text{F}_2\text{O}$: C, 58.52; H, 8.59. Found: C, 58.78; H, 8.72.

Similar reactions gave the following products.

- (1) $\text{Me}_2\text{C}(\text{OH})\text{CF}_2\text{CH}=\text{CH}_2$, $n^{25}\text{D}$ 1.3950, from 30 mmol of $\text{CH}_2=\text{CHCF}_2\text{Br}$, 42 mmol of anhydrous acetone and 25 mmol of n -butyllithium, in 41% yield. ^1H NMR($\text{CCl}_4/\text{C}_6\text{H}_6$): δ 1.17 (s, 6H, Me_2C), 1.57 (s, 1H, OH) and 5.23-6.54 (m, 3H, $\text{CH}=\text{CH}_2$).

Anal. Calcd. for $\text{C}_6\text{H}_{10}\text{F}_2\text{O}$: C, 52.93; H, 7.40. Found: C,

52.76; H, 7.52.

- (2) $\text{Et}_2\text{C}(\text{OH})\text{CF}_2\text{CH}=\text{CH}_2$, from 3 mmol of $\text{CH}_2=\text{CHCF}_2\text{Br}$, 6 mmol of Et_2CO and 3 mmol of *n*-butyllithium, in 70% yield. The spectroscopic properties of this product were in good agreement with those reported earlier.¹
- (3) $\text{cyclo-C}_6\text{H}_{10}(\text{OH})\text{CF}_2\text{CH}=\text{CH}_2$, mp 27.0-27.5°C (GLC sample), from 11.5 mmol of $\text{CH}_2=\text{CHCF}_2\text{Br}$, 20.3 mmol of cyclohexanone and 13 mmol of *n*-butyllithium, in 59% yield.
 $^1\text{H NMR}(\text{CCl}_4/\text{C}_6\text{H}_6)$: δ 0.89-1.97 (m with 2 broad s at 1.22 and 1.52 ppm, 11H, cyclohexyl H and OH); 5.21-6.44 (m, 3H, $\text{CH}=\text{CH}_2$).
 Anal. Calcd. for $\text{C}_9\text{H}_{14}\text{F}_2\text{O}$: C, 61.35; H, 8.01. Found: C, 61.44; H, 8.09.
- (4) $\text{Ph}(\text{CH}_3)\text{C}(\text{OH})\text{CF}_2\text{CH}=\text{CH}_2$, n^{25}_D 1.4955, from 10.8 mmol of $\text{CH}_2=\text{CHCF}_2\text{Br}$, 2.17 mmol of acetophenone and 12 mmol of *n*-butyllithium, in 73% yield.
 $^1\text{H NMR}(\text{CCl}_4/\text{TMS})$: δ 1.7 (s, 3H, CH_3), 2.3 (s, 1H, OH), 5.2-6.4 (m, 3H, $\text{CH}=\text{CH}_2$) and 7.4 ppm (s, 5H, Ph).
 Anal. Calcd. for $\text{C}_{11}\text{H}_{12}\text{F}_2\text{O}$: C, 66.66; H, 6.10. Found: C, 66.44; H, 6.12.
- (5) $(\text{CH}_3)_3\text{CCH}(\text{OH})\text{CF}_2\text{CH}=\text{CH}_2$, n^{25}_D 1.4116, from 9.1 mmol of $\text{CH}_2=\text{CHCF}_2\text{Br}$, 18.0 mmol of $(\text{CH}_3)_3\text{CCHO}$ and 10 mmol of *n*-butyllithium, in 95% yield.
 $^1\text{H NMR}(\text{CCl}_4/\text{C}_6\text{H}_6)$: δ 0.88 (s, 9H, CH_3), 1.60-1.86 (m, 1H, $\text{CH}(\text{OH})$), 3.05-3.67 (m, 1H, OH) and 5.22-6.55 (m, 3H, $\text{CH}=\text{CH}_2$).
 Anal. Calcd. for $\text{C}_8\text{H}_{14}\text{F}_2\text{O}$: C, 58.52; H, 8.54. Found: C, 58.42; H, 8.72.

- (6) $\text{CH}_2=\text{CHCH}(\text{OH})\text{CF}_2\text{CH}=\text{CH}_2$, n^{25}_D 1.4118, from 10.8 mmol of $\text{CH}_2=\text{CHCF}_2\text{Br}$, 30.0 mmol of $\text{CH}_2=\text{CHCHO}$ and 10 mmol of *n*-butyllithium, 20% yield.

^1H NMR($\text{CCl}_4/\text{C}_6\text{H}_6$): δ 2.10 (s, 1H, OH), 4.27 (m, 1H, $-\text{CH}-$ (OH)) and 5.10-6.55 (m, 6H, $\text{CH}=\text{CH}_2$).

Anal. Calcd. for $\text{C}_6\text{H}_8\text{F}_2\text{O}$: C, 53.73; H, 6.01. Found: C, 53.66; H, 6.25.

Also formed was $\text{CH}_2=\text{CHCH}(\text{OH})\text{C}_4\text{H}_9-\underline{n}$, n^{20}_D 1.4333 (lit.²⁵ n^{20}_D 1.4337), in 51% yield.

^1H NMR($\text{CCl}_4/\text{C}_6\text{H}_6$): δ 0.63-1.57 (m, 9H, C_4H_9), 1.70 (s, 1H, OH), 3.97 (m, 1H, $-\text{CH}(\text{OH})$) and 4.73-6.06 ppm (m, 3H, $\text{CH}=\text{CH}_2$).

- (7) $\text{PhCH}(\text{OH})\text{CF}_2\text{CH}=\text{CH}_2$, n^{25}_D 1.5008, from 6 mmol of $\text{CH}_2=\text{CHCF}_2\text{Br}$, 12 mmol of benzaldehyde and 7 mmol of *n*-butyllithium in 15% yield.

^1H NMR(CDCl_3): δ 2.4 (s, 1H, OH), 4.9 (t, $J=8\text{Hz}$, 1H, $\text{CH}-\text{CF}_2$), 5.25-6.3 (m, 3H, $\text{CH}=\text{CH}_2$) and 7.4 ppm (m, 5H, Ph).

Anal. Calcd. for $\text{C}_{10}\text{H}_{10}\text{F}_2\text{O}$: C, 65.2; H, 5.43. Found: C, 64.9; H, 5.62.

In another experiment, $\text{PhCH}(\text{OH})\text{C}_4\text{H}_9-\underline{n}$ also was identified (78%) and isolated; n^{20}_D 1.5083 (lit.²⁶ n^{20}_D 1.5078).

^1H NMR(CCl_4/TMS): δ 0.9-1.9 (m, 10H, C_4H_9 and OH), 4.6 (t, $^3J(\text{H}-\text{H})$ 7Hz, 1H, PhCH), and 7.2 ppm (s, 5H, Ph).

Reactions of gem-Difluoroallyllithium with Esters

The reaction of gem-difluoroallyllithium with methyl chloroacetate is typical.

The standard apparatus was charged with 80 ml of THF, 1.63 g

(10.7 mmol) of $\text{CH}_2=\text{CHCF}_2\text{Br}$, 3.02 g (27.8 mmol) of methyl chloroacetate, 20 ml of diethyl ether and 20 ml of pentane. This mixture was cooled to -95°C and 4.5 ml (10 mmol) of 2.3 M n-butyllithium in hexane was added dropwise, very slowly. Upon completion of the addition the reaction mixture was stirred at -95°C for 90 min and subsequently was allowed to warm slowly to room temperature. Hydrolysis with saturated aqueous NH_4Cl and work-up as in the other experiments followed. GLC analysis of the organic residue (15% QF-1 on Chromosorb W, 100°C) showed the presence of $\text{Cl-CH}_2\text{C(O)CF}_2\text{CH=CH}_2$, n^{20}_D 1.4122, in 95% yield. ^1H NMR($\text{CCl}_4/-\text{C}_6\text{H}_6$): δ 4.10 (s, 2H, ClCH_2) and 5.50-5.82 (m, 3H, CH=CH_2). IR (thin film): ν (C=O) 1752 cm^{-1} . Anal. Calcd. for $\text{C}_5\text{H}_5\text{ClF}_2\text{O}$: C, 38.86; H, 3.26. Found: C, 39.21; H, 3.50.

The following were prepared by this procedure.

- (1) $(\text{CH}_3)_2\text{CHC(O)CF}_2\text{CH=CH}_2$, n^{25}_D 1.3792, from 11.5 mmol of $\text{CH}_2=\text{CHCF}_2\text{Br}$, 30.1 mmol of $(\text{CH}_3)_2\text{CHCO}_2\text{CH}_3$ and 10 mmol of n-butyllithium, in 62% yield.
 ^1H NMR($\text{CCl}_4/\text{C}_6\text{H}_6$): δ 1.21 (d, $^3\text{J}(\text{H-H})$ 7Hz, 6H, CH_3), 3.13 (sept, $^3\text{J}(\text{H-H})$ 7Hz, 1H, Me_2CH) and 5.38-6.27 (m, 3H, CH=CH_2). IR (thin film): ν (C=O) 1740 cm^{-1} .
 Anal. Calcd. for $\text{C}_7\text{H}_{10}\text{F}_2\text{O}$: C, 56.75; H, 6.80. Found: C, 56.75; H, 6.90.
- (2) $(\text{CH}_3)_3\text{CC(O)CF}_2\text{CH=CH}_2$, n^{25}_D 1.3902, from 16.0 mmol of $\text{CH}_2=\text{CHCF}_2\text{Br}$, 20.3 mmol of $(\text{CH}_3)_3\text{CCO}_2\text{CH}_3$ and 15 mmol of n-butyllithium, in 49% yield.
 ^1H NMR($\text{CCl}_4/\text{C}_6\text{H}_6$): δ 1.30 (t, $^5\text{J}(\text{H-F})$ 0.8Hz, 9H, CH_3) and 5.36-6.13 (m, 3H, CH=CH_2). IR (thin film): ν (C=O)

1730 cm^{-1} .

Anal. Calcd. for $\text{C}_8\text{H}_{12}\text{F}_2\text{O}$: C, 59.25; H, 7.46. Found: C, 59.26; H, 7.49.

Reaction of Methyl Chloroacetate with Two Molar Equivalents of gem-Difluoroallyllithium

Using the usual procedure, 4.1 mL of 2.4M $n\text{-BuLi}$ (9.8 mmol) in hexane was added dropwise, very slowly to a solution of 4.50 g (28.7 mmol) of $\text{CH}_2=\text{CHCF}_2\text{Br}$ in 80 mL of THF, 20 mL of diethyl ether and 20 mL of pentane. The resulting mixture was stirred at -95°C for 60 min and then 4.5 mL (11 mmol) of $n\text{-BuLi}$ solution was added. After the reaction mixture had been stirred for another 60 min, 2.5 mL (20 mmol) of Me_3SiCl was added at -95°C and then the mixture was stirred and allowed to warm slowly to room temperature. After trap-to-trap distillation (70°C , 0.10 mmHg) into liquid nitrogen-cooled receiver and concentration of the volatiles at reduced pressure, GLC analysis of the liquid residue showed the presence of a major product which was identified as $\text{ClCH}_2\text{C}(\text{OMe})(\text{OSiMe}_3)\text{CF}_2\text{CH}=\text{CH}_2$ (36% yield), n_D^{20} 1.42222. ^1H NMR($\text{CCl}_4/\text{C}_6\text{H}_6$): δ 0.16 (s, 9H, SiMe_3), 3.41 (s, 3H, OMe), 3.67 (s, 2H, ClCH_2) and 5.30-6.10 ppm (m, 3H, $\text{CH}=\text{CH}_2$). Anal. Calcd. for $\text{C}_9\text{H}_{17}\text{ClF}_2\text{O}_2\text{Si}$: C, 41.77; H, 6.62. Found: C, 42.02; H, 6.65.

Acknowledgment. This work was supported in part by the U.S. Office of Naval Research.

REFERENCES AND NOTES

1. Seyferth, D.; Wursthorn, K. R. J. Organomet. Chem., 1979, 182, 455.
2. a) Seyferth, D.; Weiner, M. A. J. Am. Chem. Soc. 1962, 84, 361.
b) Schöllkopf, U., in "Houben-Weyl Methoden der Organischen Chemie", Vol. XIII/1, E. Müller, ed., Georg Thieme-Verlag, Stuttgart, 1970, pp. 130-134.
3. a) Wakefield, B. J. "The Chemistry of Organolithium Compounds", Pergamon Press, Oxford, 1974, Chapter 4.
b) Ref. 2b, pp. 148-160.
4. a) Ref. 3a, pp. 51-52.
b) Ward, H. R.; Lawler, R. G.; Loken, H. Y. J. Am. Chem. Soc. 1968, 90, 7359.
c) Ward, H. R.; Lawler, R. G.; Cooper, R. A. J. Am. Chem. Soc. 1969, 91, 746.
d) Lepley, A. R. J. Chem. Soc. Chem. Commun. 1969, 64.
e) Lepley, A. R.; Landau, R. L. J. Am. Chem. Soc. 1969, 91, 749.
5. Tarrant, P.; Lovelace, A. M. J. Am. Chem. Soc. 1954, 76, 3466.
6. Twelve times more (in terms of molar equivalents) of benzoyl peroxide was used than previous workers⁷ had used in similar Bz₂O₂-catalyzed addition of CCl₄ to ethylene.
7. Joyce, R. M.; Hanford, W. E.; Harmon, J. J. Am. Chem. Soc. 1948, 70, 2529.

8. Seyferth, D.; Wursthorn, K. R.; Lim, T.F.O.; Sepelak, D. J. J. Organomet. Chem. 1981, 205, 301.
9. a) Neugeberger, W.; Schleyer, P. V. R. J. Organomet. Chem. 1980, 198, C1.
b) Brownstein, S.; Bywater, S.; Worsfold, D. J. J. Organomet. Chem. 1980, 199, 1.
c) Bywater, S.; Patmore, D. J.; Worsfold, D. J. J. Organomet. Chem. 1977, 135, 145.
10. a) Cram, D. J., "Fundamentals of Carbanion Chemistry", Academic Press, New York, 1965, Chapter II.
b) Sheppard, W. A.; Sharts, C. M. "Organic Fluorine Chemistry", Benjamin, New York, 1969, pp.308-309.
11. Streitwieser, A.; Mares, F. J. Am. Chem. Soc. 1968, 90, 2444.
12. Hine, J.; Mahone, L. G.; Liotta, C. L. J. Am. Chem. Soc. 1967, 89, 5911.
13. In support of this idea, we note that CNDO/2MO calculations have indicated that in $\text{Li}[\text{CF}_2\text{CHCH}_2] \cdot 2\text{Me}_2\text{O}$ the total charge density at the CH_2 terminus is much greater than at the CF_2 terminus: Prof. J. F. Sebastian (Miami University), private communication, October 1974.
14. Seyferth, D.; Murphy, G. J.; Mauzé, B. J. Am. Chem. Soc. 1977, 99, 5317.
15. Adolph, H. G.; Kamlet, M. J. J. Am. Chem. Soc. 1966, 88, 4761.
16. Mauzé, B. J. Organomet. Chem. 1979, 170, 265.
17. Doucoure, A.; Mauzé, B.; Miginiac, L. J. Organomet. Chem. 1982, 236, 139.

18. Seyferth, D.; Mammarella, R. E. J. Organomet. Chem. 1978, 156, 279.
19. Seyferth, D.; Mammarella, R. E. J. Organomet. Chem. 1979, 177, 53.
20. a) van Dongen, J. P.; van Dijkman, H. W.; de Bie, M. J. Recl. Trav. Chim. Pays-Bas 1974, 93, 29.
b) Bywater, S.; Lachance, P.; Worsfold, D. J. J. Phys. Chem. 1975, 79, 2148.
c) Bywater, S.; Worsfold, D. J. J. Organomet. Chem. 1978, 159, 229.
21. Courtois, G.; Miginiac, L. J. Organomet. Chem. 1974, 69, 1.
22. a) Ref. 10b, Chapter 9, Sect. 3.
b) Schlosser, M. Tetrahedron 1978, 34, 3.
c) Walsh, C. T., Advan. Enzymol. 1983, 54, in press.
23. Kofron, W. G.; Baclawski, L. M. J. Org. Chem. 1976, 41, 1879.
24. Walters, R. J.; Tract, J. H.; Weinburger, E. B.; Rodgers, J. K. Chem. Eng. Prog. 1954, 50, 511.
25. Grayson, J. T.; Greenlee, K. W.; Derfer, J. M. Boord, C. E. J. Org. Chem. 1955, 20, 275.
26. Ipatieff, V. N.; Haensel, V. J. Am. Chem. Soc. 1942, 64, 520.

TECHNICAL REPORT DISTRIBUTION LIST, GEN

	<u>No.</u> <u>Copies</u>	<u>Co</u>
Office of Naval Research Attn: Code 413 800 North Quincy Street Arlington, Virginia 22217	2	
ONR Pasadena Detachment Attn: Dr. R. J. Marcus 1030 East Green Street Pasadena, California 91106	1	
Commander, Naval Air Systems Command Attn: Code 310C (H. Rosenwasser) Department of the Navy Washington, D.C. 20360	1	
Defense Technical Information Center Building 5, Cameron Station Alexandria, Virginia 22314	12	
Dr. Fred Saalfeld Chemistry Division, Code 6100 Naval Research Laboratory Washington, D.C. 20375	1	
U.S. Army Research Office Attn: CRD-AA-IP P. O. Box 12211 Research Triangle Park, N.C. 27709	1	
Mr. Vincent Schaper DTNSRDC Code 2803 Annapolis, Maryland 21402	1	
Naval Ocean Systems Center Attn: Dr. S. Yamamoto Marine Sciences Division San Diego, California 91232	1	
		Naval Ocean Systems Center Attn: Mr. Joe McCartney San Diego, California 92152
		Naval Weapons Center Attn: Dr. A. B. Amster, Chemistry Division China Lake, California 93555
		Naval Civil Engineering Laboratory Attn: Dr. R. W. Drisko Port Hueneme, California 93401
		Dean William Tolles Naval Postgraduate School Monterey, California 93940
		Scientific Advisor Commandant of the Marine Corps (Code RD-1) Washington, D.C. 20380
		Naval Ship Research and Development Center Attn: Dr. G. Bosmajian, Applied Chemistry Division Annapolis, Maryland 21401
		Mr. John Boyle Materials Branch Naval Ship Engineering Center Philadelphia, Pennsylvania 19112
		Mr. A. M. Anzalone Administrative Librarian PLASTEC/ARRADCOM Bldg 3401 Dover, New Jersey 07801

TECHNICAL REPORT DISTRIBUTION LIST, 356B

	<u>No.</u> <u>Copies</u>	<u>N</u> <u>Cop</u>
Dr. C. L. Shilling ^c Union Carbide Corporation Chemical and Plastics Tarrytown Technical Center Tarrytown, New York	1	
Dr. R. Soulen Contract Research Department Pennwalt Corporation 900 First Avenue King of Prussia, Pennsylvania 19406	1	
Dr. A. G. MacDiarmid University of Pennsylvania Department of Chemistry Philadelphia, Pennsylvania 19174	1	
Dr. H. Allcock Pennsylvania State University Department of Chemistry University Park, Pennsylvania 16802	1	
Dr. M. Kenney Case-Western University Department of Chemistry Cleveland, Ohio 44106	1	
Dr. R. Lenz University of Massachusetts Department of Chemistry Amherst, Massachusetts 01002	1	
DR. M. David Curtis University of Michigan Department of Chemistry Ann Arbor, Michigan 48105	1	
NASA-Lewis Research Center Attn: Dr. T. T. Serafini, MS 49-1 21000 Brookpark Road Cleveland, Ohio 44135	1	
Dr. J. Griffith Naval Research Laboratory Chemistry Section, Code 6120 Washington, D.C. 20375	1	
Dr. G. Goodman Globe-Union Incorporated 5757 North Green Bay Avenue Milwaukee, Wisconsin 53201		
Dr. E. Fischer, Code 2853 Naval Ship Research and Development Center Annapolis Division Annapolis, Maryland 21402		
Dr. Martin H. Kaufman Code 38506 Naval Weapons Center China Lake, California 93555		
Dr. C. Allen University of Vermont Department of Chemistry Burlington, Vermont 05401		
Professor R. Drago Department of Chemistry University of Florida Gainesville, FL 32611		
Dr. D. L. Venezky Code 6130 Naval Research Laboratory Washington, D.C. 20375		
COL R. W. Bowles, Code 100M Office of Naval Research 800 N. Quincy Street Arlington, Virginia 22217		
Professor T. Katz Department of Chemistry Columbia University New York, New York 10027		
Professor James Chien Department of Chemistry University of Massachusetts Amherst, Massachusetts 01002		

TECHNICAL REPORT DISTRIBUTION LIST, 356B

	<u>No. Copies</u>
Professor Malcolm B. Polk Department of Chemistry Atlanta University Atlanta, Georgia 30314	1
Dr. G. Bryan Street IBM Research Laboratory, K32/281 San Jose, California 95193	1
Professor Michael Moran Department of Chemistry West Chester State College West Chester, Pennsylvania 19401	1
Dr. K. Paciorek Ultrasystems, Inc. P. O. Box 19605 Irvine, California 92715	1
Dr. D. B. Cotts SRI International 333 Ravenswood Avenue Menlo Park, California 94025	1
Professor D. Seyferth Department of Chemistry Massachusetts Institute of Technology Cambridge, Massachusetts 02139	1
Dr. Kurt Baum Fluorochem, Inc. 680 S. Aycn Avenue Azusa, California 91702	1

